

A Test of CPT Symmetry in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ Decays^{*}

XING Zhi-Zhong^{1,2;1)}

¹ (Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China)

² (Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences, Beijing 100049, China)

Abstract I show that the CP-violating asymmetry in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ decays differs from that in $K_L \rightarrow \pi^+\pi^-$, $K_L \rightarrow \pi^0\pi^0$ or the semileptonic K_L transitions, if there exists CPT violation in K^0 - $\bar{K}^0$ mixing. A delicate measurement of this difference at a super flavor factory (e.g., the ϕ factory) will provide us with a robust test of CPT symmetry in the neutral kaon system.

Key words K^0 - $\bar{K}^0$ mixing, CPT violation

PACS 11.30. Er, 13.25. Es, 14.40. Aq

1 The motivation

The CPT theorem claims that a Lorentz-invariant local quantum field theory with a Hermitian Hamiltonian must have CPT symmetry [1]. It is so far so good, because there is no convincing experimental hint at CPT violation [2]. The breaking of CPT symmetry, as expected in some “exotic” scenarios of new physics beyond the standard model (e.g., string theory) [3], would be a big deal. In any case, much more experimental tests of this theorem are desirable.

The K^0 - $\bar{K}^0$ mixing system has been playing an important role in particle physics for testing fundamental symmetries (such as CP, T and CPT) and examining conservation laws (such as $\Delta S = \Delta Q$). The existing experimental evidence for CPT invariance in the mixing and decays of neutral kaon mesons remains rather poor [2]: it is not excluded that the strength of CPT-violating interactions could be as large as about ten percentage of that of CP-violating interactions. This unsatisfactory situation will be improved in the near future, in particular after a variety of more delicate measurements are carried out at a super flavor factory [4] (e.g., the ϕ factory [5]).

There are several possibilities of probing CPT violation in K^0 - $\bar{K}^0$ mixing with the decays of K_S and K_L mesons into the two-pion and (or) the semileptonic states [2]. A different approach towards testing CPT symmetry, with the help of neutral kaon de-

cays into the three-pion states, has also been pointed out in Ref. [6]. The idea is simply that the CP-violating effect induced by K^0 - $\bar{K}^0$ mixing in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ transitions should not be identical to that in $K_L \rightarrow \pi^+\pi^-$, $K_L \rightarrow \pi^0\pi^0$ or the semileptonic K_L decays, if CPT symmetry is broken. Thus a careful comparison between these two types of CP-violating effects may provide us with a robust test of CPT invariance in K^0 - $\bar{K}^0$ mixing.

An unfortunate fact is that no attention has so far been paid to the method advocated in Ref. [6]. In this talk, which is more or less an advertisement, I shall explain why a test of CPT symmetry is possible by measuring the time-dependent CP-violating asymmetry between $K^0(t) \rightarrow \pi^+\pi^-\pi^0$ and $\bar{K}^0(t) \rightarrow \pi^+\pi^-\pi^0$ decays. My result is hopefully useful for the upcoming experiments of kaon physics.

2 The idea

Let me outline the main idea. The mass eigenstates of K^0 and $\bar{K}^0$ can in general be written as

$$\begin{aligned} |K_S\rangle &= \frac{1}{\sqrt{|p_1|^2 + |q_1|^2}} (p_1|K^0\rangle + q_1|\bar{K}^0\rangle), \\ |K_L\rangle &= \frac{1}{\sqrt{|p_2|^2 + |q_2|^2}} (p_2|K^0\rangle - q_2|\bar{K}^0\rangle), \end{aligned} \quad (1)$$

in which p_i and q_i (for $i = 1, 2$) are complex mixing parameters. Note that $p_1 = p_2$ and $q_1 = q_2$ follow

^{*} Supported in part by the National Natural Science Foundation of China (10425522 and 10875131) and in part by the Ministry of Science and Technology of China (2009CB825207).

1) E-mail: xingzz@ihep.ac.cn

from CPT invariance [7]. The traditional characteristic quantities of CP violation in the K^0 - $\bar{K}^0$ mixing system [2], η_{+-} , η_{00} and δ_L , are all related to K_L decays and thus the (p_2, q_2) parameters. For example,

$$\delta_L \equiv \frac{|p_2|^2 - |q_2|^2}{|p_2|^2 + |q_2|^2} \quad (2)$$

in the absence of $\Delta S = -\Delta Q$ interactions. A measurement of CP violation associated with

$$\delta_S = \frac{|p_1|^2 - |q_1|^2}{|p_1|^2 + |q_1|^2} \quad (3)$$

has been assumed to be extremely difficult, if not impossible, due to the rapid decay of the K_S meson to the two-pion state or the semileptonic state. Nevertheless, I shall show that δ_S can be measured from the rate asymmetry of K^0 and $\bar{K}^0$ mesons decaying into the three-pion state $\pi^+\pi^-\pi^0$. The difference between δ_S and δ_L signifies CPT violation in K^0 - $\bar{K}^0$ mixing. This point can be seen more clearly if one adopts the popular (ϵ, δ) parameters to describe CP- and CPT-violating effects in the K^0 - $\bar{K}^0$ mixing system [2]:

$$\begin{aligned} p_1 &= 1 + \epsilon + \delta, \\ p_2 &= 1 + \epsilon - \delta, \\ q_1 &= 1 - \epsilon - \delta, \\ q_2 &= 1 - \epsilon + \delta. \end{aligned} \quad (4)$$

Then

$$\begin{aligned} \delta_L &= 2(\text{Re}\epsilon - \text{Re}\delta), \\ \delta_S &= 2(\text{Re}\epsilon + \text{Re}\delta). \end{aligned} \quad (5)$$

It turns out that $\delta_S - \delta_L = 4\text{Re}\delta$ is a clear signature of CPT violation [6].

Let me quote two typical experimental constraints on the CPT-violating parameter δ in K^0 - $\bar{K}^0$ mixing: $\text{Re}\delta = (2.9 \pm 2.6_{\text{stat}} \pm 0.6_{\text{syst}}) \times 10^{-4}$ obtained by the

CPLEAR Collaboration [8] and $\text{Im}\delta = (0.4 \pm 2.1) \times 10^{-5}$ obtained by the KLOE Collaboration [9]. A systematic analysis of the CP- and CPT-violating parameter space has already been done by the Particle Data Group in Ref. [2].

3 The approach

The CP eigenvalue for the $\pi^+\pi^-\pi^0$ final state is given by $(-1)^{l+1}$, where l is the relative angular momentum between π^+ and π^- . Since the sum of the masses of three pions is close to the kaon mass, the pions have quite low kinetic energy $E_{\text{CM}}(\pi)$ in the kaon rest-frame, and the states with $l > 0$ are suppressed by the centrifugal barrier [10]. Thus the K_L meson decays dominantly into the kinematics-favored ($l = 0$) and CP-allowed (CP = -1) $\pi^+\pi^-\pi^0$ state. The decay amplitude of $K_S \rightarrow \pi^+\pi^-\pi^0$ consists of both the kinematics-suppressed ($l = 1$) but CP-allowed (CP=+1) component, and the kinematics-favored ($l = 0$) but CP-forbidden (CP=-1) component. This implies an interesting Dalitz-plot distribution for the $K_S \rightarrow \pi^+\pi^-\pi^0$ transition: it is symmetric with respect to π^+ and π^- for the CP-violating amplitude, but anti-symmetric for the CP-conserving amplitude. Let the ratio of K_S and K_L decay amplitudes be

$$\eta_{+-0} = \frac{A(K_S \rightarrow \pi^+\pi^-\pi^0)}{A(K_L \rightarrow \pi^+\pi^-\pi^0)}. \quad (6)$$

It is clear that η_{+-0} depends only upon the CP-violating component of $A(K_S \rightarrow \pi^+\pi^-\pi^0)$, if data are integrated over the whole Dalitz plot [10, 11]. The time-dependent rates for the initially pure K^0 and $\bar{K}^0$ states decaying into $\pi^+\pi^-\pi^0$, denoted by $\mathcal{R}(t)$ and $\bar{\mathcal{R}}(t)$ respectively, can be calculated with the help of Eqs. (1) and (6). I arrive at [6]

$$\begin{aligned} \mathcal{R}(t) &\propto [|q_1|^2 + |q_2|^2 |\eta_{+-0}|^2 e^{-\Delta\Gamma t} + 2\text{Re}(q_1^* q_2 \eta_{+-0} e^{i\Delta m t}) e^{-\Delta\Gamma t/2}], \\ \bar{\mathcal{R}}(t) &\propto [|p_1|^2 + |p_2|^2 |\eta_{+-0}|^2 e^{-\Delta\Gamma t} - 2\text{Re}(p_1^* p_2 \eta_{+-0} e^{i\Delta m t}) e^{-\Delta\Gamma t/2}], \end{aligned} \quad (7)$$

where $\Delta m > 0$ and $\Delta\Gamma > 0$ denote the mass difference and the width difference of K_S and K_L mesons,

respectively. To a good degree of accuracy, I obtain the following CP-violating asymmetry:

$$\begin{aligned} \mathcal{A}(t) \equiv \frac{\bar{\mathcal{R}}(t) - \mathcal{R}(t)}{\bar{\mathcal{R}}(t) + \mathcal{R}(t)} &= \delta_S - 2e^{-\Delta\Gamma t/2} [\text{Re}\eta_{+-0} \cos(\Delta m t) - \text{Im}\eta_{+-0} \sin(\Delta m t)] \xi \\ &\quad - 2e^{-\Delta\Gamma t/2} [\text{Re}\eta_{+-0} \sin(\Delta m t) + \text{Im}\eta_{+-0} \cos(\Delta m t)] \zeta, \end{aligned} \quad (8)$$

in which

$$\begin{aligned}\xi &= \frac{\text{Re}(p_1 p_2^* + q_1 q_2^*)}{|p_1|^2 + |q_1|^2} = 1 + \mathcal{O}(|\epsilon|^2) + \mathcal{O}(|\delta|^2) + \mathcal{O}(\text{Re}(\epsilon\delta^*)), \\ \zeta &= \frac{\text{Im}(p_1 p_2^* + q_1 q_2^*)}{|p_1|^2 + |q_1|^2} = \mathcal{O}(\text{Im}(\epsilon\delta^*)).\end{aligned}\quad (9)$$

It is obvious that δ_S can be determined through the measurement of $\mathcal{A}(t)$. In particular, the relationship $\lim_{t \rightarrow \infty} \mathcal{A}(t) = \delta_S$ holds.

As I have emphasized, the difference between δ_S and δ_L hints at CPT violation in K^0 - $\bar{K}^0$ mixing. If

$|\text{Re}\delta|/\text{Re}\epsilon \sim 0.1$, then the difference $\delta_S - \delta_L = 4\text{Re}\delta$ can be as large as $0.4\text{Re}\epsilon \sim 6.6 \times 10^{-4}$ in magnitude, where the experimental value $\text{Re}\epsilon \approx 1.65 \times 10^{-3}$ has been used [2]. Since both ϵ and δ are small quantities, it turns out that $\xi \approx 1$ and $\zeta \approx 0$ are good approximations. Eq. (8) is therefore simplified to

$$\mathcal{A}(t) = \delta_S - 2e^{-\Delta\Gamma t/2} [\text{Re}\eta_{+-0} \cos(\Delta m t) - \text{Im}\eta_{+-0} \sin(\Delta m t)]. \quad (10)$$

In the neglect of CPT violation, namely, $\delta_S = 2\text{Re}\epsilon$, Eq. (10) can simply reproduce the result obtained in Ref. [10]. For illustration, I plot the behavior of $\mathcal{A}(t)$ in Fig. 1, in which $\delta_S = 3 \times 10^{-3}$ and $|\eta_{+-0}| = 5 \times 10^{-3}$ have typically been input. One may observe that $\mathcal{A}(t)$ approaches δ_S for $t \geq 5\tau_S$ and reaches δ_S if $t \geq 10\tau_S$, where τ_S is the mean lifetime of the K_S meson. This implies a certain feasibility to determine δ_S from the time-dependent CP-violating asymmetry $\mathcal{A}(t)$.




Fig. 1. An illustrative plot for the CP-violating asymmetry $\mathcal{A}(t)$ with the typical inputs $\delta_S = 3 \times 10^{-3}$ and $|\eta_{+-0}| = 5 \times 10^{-3}$ [6].

4 The discussion

In the above analysis I have taken an integration over the whole Dalitz plot, such that η_{+-0} solely contains the CP-violating part of $A(K_S \rightarrow \pi^+ \pi^- \pi^0)$. To look at the CP-conserving component of $A(K_S \rightarrow \pi^+ \pi^- \pi^0)$, one may study the phase-space regions

$E_{\text{CM}}(\pi^+) > E_{\text{CM}}(\pi^-)$ and $E_{\text{CM}}(\pi^+) < E_{\text{CM}}(\pi^-)$ separately [10]. In this case the corresponding CP-violating asymmetries between $\bar{\mathcal{R}}(t)$ and $\mathcal{R}(t)$ take the same form as $\mathcal{A}(t)$ in Eq. (8) or Eq. (10), but η_{+-0} should be replaced by $(\eta_{+-0} \pm \lambda)$, where λ denotes the CP-conserving contribution to the ratio of K_S and K_L decay amplitudes [10]. Certainly, the CP-violating parameter δ_S can still be determined from measuring the time dependence of the relevant decay rate asymmetries.

An accurate measurement of δ_S from K^0 vs $\bar{K}^0 \rightarrow \pi^+ \pi^- \pi^0$ should be feasible at the ϕ factory, where a huge amount of $K^0 \bar{K}^0$ events can be coherently produced [5]. Choosing the semileptonic decay of one kaon to tag the flavor of the other kaon decaying into $\pi^+ \pi^- \pi^0$ on the ϕ resonance, one should be able to construct the time-dependent rate asymmetry between $K^0(t) \rightarrow \pi^+ \pi^- \pi^0$ and $\bar{K}^0(t) \rightarrow \pi^+ \pi^- \pi^0$ decays in a way similar to Eq. (8). It is also expected that other super flavor factories may measure δ_S and δ_L to a good degree of accuracy.

Note that Lorentz invariance has been taken for granted in what I have discussed. As pointed out by Greenberg [12], “If CPT invariance is violated in an interacting quantum field theory, then that theory also violates Lorentz invariance”. In my discussions, the dependence of the CPT-violating parameter δ on the sidereal time should in general be considered, since CPT violation may simultaneously imply a violation of Lorentz symmetry in the neutral kaon system. For simplicity, here I take δ to be a constant by assuming that the boost parameters of both K^0 and $\bar{K}^0$ are small and the corresponding Lorentz-

violating effect is rotationally invariant in the laboratory frame [13]. In this approximation, my results are essentially valid as the averages over the sidereal time, such that the effect of Lorentz violation due to the direction of motion is negligible.

Finally, I like to mention that different approaches have been discussed to test CPT symmetry in D^0 - $\bar{D}^0$, B_d^0 - $\bar{B}_d^0$ or B_s^0 - $\bar{B}_s^0$ mixing [14]. The idea presented here cannot directly be applied to those heavy neutral-meson systems. In this sense, it represents a unique way applicable in the K^0 - $\bar{K}^0$ mixing system to test the CPT theorem.

5 The conclusion

To conclude, the CP-violating effect induced by K^0 - $\bar{K}^0$ mixing in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ decays is possible to deviate to some extent from that in $K_L \rightarrow \pi\pi$ or the semileptonic K_L transitions due to the violation of CPT symmetry. Measuring or constraining this tiny difference may serve as a robust test of CPT invariance in the neutral kaon system.

I would like to thank Changzheng Yuan and the organizing committee of PHIPSI09 for giving me this opportunity to present an “old” idea for the future.

References

- 1 J. Schwinger, Phys. Rev. **82**, 914 (1951); G. Lüders, Kgl. Dansk. Vidensk. Selsk. Mat. Fys. Medd. **28**, 1 (1954); W. Pauli, in *Niels Bohr and the Development of Physics* (Pergamon Press, New York, 1955), p. 30.
- 2 Particle Data Group, C. Amsler *et al.*, Phys. Lett. B **667**, 1 (2008).
- 3 V.A. Kostelecky, hep-ph/0104227.
- 4 D. Hitlin, in these proceedings.
- 5 *The DAΦNE Physics Handbook*, edited by L. Maiani, G. Pancheri, and N. Paver (Frascati, 1992).
- 6 Z.Z. Xing, Phys. Rev. D **62**, 097901 (2000).
- 7 T.D. Lee and C.S. Wu, Ann. Rev. Nucl. Part. Sci. **16**, 511 (1966).
- 8 CPLEAR Collaboration, A. Angelopoulos *et al.*, Phys. Lett. B **444**, 52 (1998).
- 9 KLOE Collaboration, F. Ambrosino *et al.*, JHEP **0612**, 011 (2006).
- 10 CPLEAR Collaboration, R. Adler *et al.*, Phys. Lett. B **364**, 239 (1995); **370**, 167 (1996); **374**, 313 (1996); **407**, 193 (1997).
- 11 T. Nakada and L. Wolfenstein, Eur. Phys. J. C **3**, 457 (1998).
- 12 O.W. Greenberg, Phys. Rev. Lett. **89**, 231602 (2002).
- 13 V.A. Kostelecky, Phys. Rev. Lett. **80**, 1818 (1998); Phys. Rev. D **64**, 076001 (2001).
- 14 See, e.g., Z.Z. Xing, Phys. Rev. D **50**, 2957 (1994); Phys. Rev. D **55**, 196 (1997); Phys. Lett. B **450**, 202 (1999); Phys. Lett. B **487**, 327 (2000); P. Ren and Z.Z. Xing, Phys. Rev. D **76**, 116001 (2007).